Corrected Amendment and Response dated August 2, 2004

Reply to Notice of Non-compliant Amendment dated July 2, 2004

Page 3 of 14

Amendments to Claims:

Claims 23-31 and 33 are hereby withdrawn. Claims 1, 3, 14-18, and 22 are herby amended as shown in the listing below. This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

- Claim 1 (currently amended) A method for detecting an unbound form of a first member of a binding pair, the binding pair comprising a first and second member, each member bindable to the other, the method comprising the steps of:
 - (a) providing a first particle bound to the second member;
 - (b) reacting the first particle bound to the second member with a sample, thereby forming a first complex between the second member bound to the first particle and unbound first member present in said sample;
 - (c) providing a second particle bound to a third member, the third member being different from the second member and being capable of binding to the first member;
 - (d) reacting the second particle bound to the third member to-with the sample, thereby forming a second complex between the third member bound to the second particle and the first complex; and
 - (e) detecting any second complex formed by determining turbidity or agglutination.
- Claim 2 (original) The method of claim 1, wherein the third member is an antibody which specifically binds to the first member.

Corrected Amendment and Response dated August 2, 2004

Reply to Notice of Non-compliant Amendment dated July 2, 2004

Page 4 of 14

- Claim 3 (currently amended) The method of claim 1, wherein <u>at least one of</u> the first and/or second particle <u>is comprises</u> latex.
- Claim 4 (original) The method of claim 1, wherein the second complex is detected by measuring an increase in the turbidity of the sample.
- Claim 5 (original) The method of claim 1, wherein steps (a) through (d) are performed sequentially.
- Claim 6 (original) The method of claim 1, wherein steps (a) through (d) are performed simultaneously.
- Claim 7 (original) The method of claim 1, wherein the amount of second complex formed is quantitated.
- Claim 8 (original) The method of claim 1, wherein the first member is protein S.
- Claim 9 (original) The method of claim 1, wherein the second member is C4b-binding protein (C4BP).
- Claim 10 (original) The method of claim 1, wherein the sample is selected from the group consisting of blood, plasma, serum, saliva, CSF, urine, culture media, a cell suspension, a buffer and an artificially prepared fluid containing the first member.
- Claim 11 (original) The method of claim 1, wherein the second member binds to the first member at a single binding site.
- Claim 12 (original) The method of claim 11, wherein the third member binds to the first member at a single binding site which is different from the single binding site to which the second member binds.

Corrected Amendment and Response dated August 2, 2004

Reply to Notice of Non-compliant Amendment dated July 2, 2004

Page 5 of 14

- Claim 13 (original) The method of claim 1, wherein step (b) is performed within 0 to about 180 seconds.
- Claim 14 (currently amended) The method of claim 1, wherein the molar ratio of third member to and the second member are in a molar ratio of between about 2 and 20.
- Claim 15 (currently amended) The method of claim 1, wherein the molar ratio of the third member to and the second member are in a molar ratio of the between about 5 and 10.
- Claim 16 (currently amended) The method of claim 1, wherein the amount of third member is present in an amount that is higher than thean amount of the free first member is in the sample.
- Claim 17 (currently amended) The method of claim 1, wherein the molar ratio of third member and the free first member in the sample are in a molar ration of the between about 10 and 40 times the amount of free first member in the sample.
- Claim 18 (currently amended) A composition for detecting an unbound form of a first member of a binding pair, the binding pair comprising a first and <u>a</u> second member, each member bindable to the other, the composition comprising:
 - a first particle bound to the second member;
 - a second particle bound to a third member, the third member being different from the second member and capable of binding to the first member at a binding site different from the second member, wherein the first member comprises protein S and the second member comprises C4BP.
- Claim 19 (original) The composition of claim 18, wherein the first member is protein S and the second member is C4BP.

Corrected Amendment and Response dated August 2, 2004

Reply to Notice of Non-compliant Amendment dated July 2, 2004

Page 6 of 14

- Claim 20 (original) The composition of claim 18, wherein the third member is an antibody and the second member is not an antibody.
- Claim 21 (original) The composition of claim 18, wherein the second member comprises a single binding site for the first member.
- Claim 22 (currently amended) The composition of claim 21, wherein the third member binds to the first member at a single binding site which is different from the single binding site to which the second member binds.
- Claim 23 (withdrawn) A method for detecting an unbound form of a first member of a binding pair, the binding pair comprising a first and second member, each member bindable to the other, the method comprising the steps of:
 - (a) providing a first particle bound to the second member;
 - (b) reacting the first particle bound to the second member with a sample, thereby forming a first complex between the second member bound to the first particle and unbound first member present in said sample;
 - (c) providing a second particle bound to the first member;
 - (d) reacting the second particle bound to the first member with the sample, thereby forming a second complex between second particle bound to the first member and first particle bound to second member which is not already bound to the first member; and
 - (e) detecting any second complex formed, wherein the amount of second complex formed is inversely proportional to the amount of unbound first member is the sample.
- Claim 24 (withdrawn) The method of claim 23, wherein the first and/or second particle is latex.

Corrected Amendment and Response dated August 2, 2004

Reply to Notice of Non-compliant Amendment dated July 2, 2004

Page 7 of 14

- Claim 25 (withdrawn) The method of claim 23, wherein the second complex is detected by measuring an increase in the turbidity of the sample.
- Claim 26 (withdrawn) The method of claim 23, wherein the amount of second complex formed is quantitated.
- Claim 27 (withdrawn) The method of claim 23, wherein the first member is protein S.
- Claim 28 (withdrawn) The method of claim 23, wherein the second member is C4BP.
- Claim 29 (withdrawn) The method of claim 23 wherein the sample is selected from the group consisting of blood, plasma, serum, or an artificially prepared buffer containing the first member.
- Claim 30 (withdrawn) A composition for detecting an unbound form of a first member of a binding pair comprising a first and second member, each member bindable to the other, the composition comprising:
 - a first particle bound to the second member; and
 - a second particle bound to the first member.
- Claim 31 (withdrawn) The composition of claim 30, wherein the first member is protein S and the second member is C4BP.
- Claim 32 (original) A method for diagnosing thrombophilia comprising performing the method of claim 8, and further comprising comparing the amount of second complex formed to the amount of second complex formed in a sample derived from an individual without thrombophilia.
- Claim 33 (withdrawn) A method for diagnosing thrombophilia comprising performing the method of claim 27, and further comprising comparing the amount of second complex

Application Serial No.: 10/022,481

<u>Corrected</u> Amendment and Response dated August 2, 2004

Reply to Notice of Non-compliant Amendment dated July 2, 2004

Page 8 of 14

formed to the amount of second complex formed in a sample derived from an individual without thrombophilia.